

GUIDELINE

Polycythaemia

Scope (Staff):	Nursing and Medical Staff
Scope (Area):	NICU KEMH, NICU PCH, NETS WA

Child Safe Organisation Statement of Commitment

CAHS commits to being a child safe organisation by applying the National Principles for Child Safe Organisations. This is a commitment to a strong culture supported by robust policies and procedures to reduce the likelihood of harm to children and young people.

This document should be read in conjunction with this disclaimer

Aim

To outline the causes and management of neonatal polycythaemia

Risk

High viscosity due to polycythaemia can lead to decreased perfusion resulting in multiorgan dysfunction in newborn babies

Background

Polycythaemia is defined as a venous haematocrit **(HCT) > 65%.** Incidence in healthy neonates is 0.4 to 5%. Capillary HCT is generally higher than venous HCT which in turn is higher than 'central' HCT (from umbilical vein). Venous samples are preferred over heel prick capillary samples while managing neonates with polycythaemia.

The haematocrit increases after birth, peaks at 2-6 hours of life, then drops slowly to cord blood values at 18 hours, thereafter it stays relatively stable. Polycythaemia occurs as a result of active erythropoiesis or passive transfusion.

Conditions that predispose newborn infants to polycythaemia include the following:

- Delayed cord clamping
- Twin to twin transfusions (recipient twin)
- Maternal foetal transfusion
- Prenatal asphyxia
- Intrauterine hypoxia, e.g. SGA (utero-placental insufficiency)

- Maternal diabetes
- Maternal hypertension
- Maternal smoking
- Rare conditions: Beckwith-Wiedemann and Trisomies 13, 18 and 21.

Clinical Features

- Lethargy and poor feeding
- RDS (pulmonary capillary sludging), cyanosis, failure of / delayed transition
- CNS depression ~Tremors, jitteriness, seizures, coma
- Hypoglycaemia (12% to 40%)
- Hypocalcaemia
- Poor renal function (renal vein thrombosis)
- Jaundice
- NEC (mesenteric hypoxia)
- Cardiac symptoms such as tachypnoea, cyanosis, tachycardia, cardiomegaly in up to 50% of plethoric infants
- Abnormal coagulation profile, thrombocytopenia

Treatment

Management of polycythaemia in neonates is controversial (Partial exchange transfusion vs. conservative management).

Conservative management is with liberalisation of IV fluids but watch for hyponatremia.

- Partial dilution exchange transfusion (PET) with normal saline reverses the physiological abnormalities and ameliorates symptoms. It also improves cerebral blood flow and hemodynamic parameters but doesn't improve long term outcomes. PET may be associated with increased risk of NEC (*Mimouni 2011*). PET is recommended for symptomatic infants with polycythaemia and should be performed as early as possible under intensive monitoring. Usually babies become symptomatic when HCT reaches 70% or above. Occasionally, PET may be required even when HCT is 65% if baby is symptomatic.
 - The goal is to decrease the haematocrit to 55%.
 - The following formula is used to calculate the partial exchange volume ((*Mimouni 2011*).

Volume to be exchanged [Wt (kg) x 80 x (HCT of patient- Desired HCT)] ÷ HCT of patient	
--	--

• Example: 3 kg baby who has haematocrit of 70%

- Desired haematocrit=55%¹
- Volume to be exchanged (ml)= [3x80 x (70-55)] ÷70 = 51 ml
- Aliquots should not exceed 5 mL/kg and should be delivered or removed over 2-3 minutes.
- **Other management:** Carefully monitor babies for hypoglycaemia and thrombocytopenia and treat where indicated.

Related CAHS internal policies, procedures and guidelines

Neonatology Guideline

• Exchange Transfusion

References and related external legislation, policies, and guidelines

- 1. Mimouni FB et al. Neonatal polycythaemia: critical review and a consensus statement of the Israeli Neonatology Association. Acta Paediatr. 2011 Oct;100(10):1290-6.
- 2. Ozek E et al. Partial exchange transfusion to prevent neurodevelopmental disability in infants with polycythemia. Cochrane Database Syst Rev. 2010 Jan 20;(1):CD005089.
- 3. Dempsey EM et al. Crystalloid or colloid for partial exchange transfusion in neonatal polycythemia: a systematic review and meta-analysis. Acta Paediatr. 2005 Nov;94(11):1650-5
- 4. Hopfeld-Fogel A et al. Neonatal Polycythemia and Hypoglycemia in Newborns: Are They Related? Am J Perinatol. 2021 Jul;38(9):930-934.
- 5. Vlug RD et al. Thrombocytopenia in neonates with polycythemia: incidence, risk factors and

clinical outcome. Expert Rev Hematol. 2015 Feb;8(1):123-9.

- 6. Morag I et al. Restrictive management of neonatal polycythemia. Am J Perinatol. 2011 Oct;28(9):677-82.
- 7. Hopewell B et al. Partial exchange transfusion for polycythemia hyperviscosity syndrome. Am J Perinatol. 2011 Aug;28(7):557-64.
- 8. Uslu S et al. The evaluation of polycythemic newborns: efficacy of partial exchange transfusion. J Matern Fetal Neonatal Med. 2011 Dec;24(12):1492-7.
- 9. Juan I. Remon et al. Polycythemia in the Newborn. Neoreviews 2011;12;e20.

This document can be made available in alternative formats on request.

Document Owner:	Neonatology				
Reviewer / Team:	Neonatology				
Date First Issued:	June 2006	Last Reviewed:	October 2021		
Amendment Dates:		Next Review Date:	21 st October 2024		
Approved by:	Neonatal Coordinating Group	Date:	21 st October 2021		
Endorsed by:	Neonatal Coordinating Group	Date:			
Standards Applicable:	NSQHS Standards: 90 Child Safe Standards: 1,10				
Printed or personally saved electronic copies of this document are considered uncontrolled					
Healthy kids, healthy communities Compassion Excellence Collaboration Accountability Equity Respect Neonatology Community Health Mental Health Perth Children's Hospital					